


Guidelines for modelling radionuclide environmental transport, fate, and exposure associated with the normal operation of nuclear facilities



Legal Notice for Standards

Canadian Standards Association (operating as “CSA Group”) develops standards through a consensus standards development process approved by the Standards Council of Canada. This process brings together volunteers representing varied viewpoints and interests to achieve consensus and develop a standard. Although CSA Group administers the process and establishes rules to promote fairness in achieving consensus, it does not independently test, evaluate, or verify the content of standards.

Disclaimer and exclusion of liability

This document is provided without any representations, warranties, or conditions of any kind, express or implied, including, without limitation, implied warranties or conditions concerning this document’s fitness for a particular purpose or use, its merchantability, or its non-infringement of any third party’s intellectual property rights. CSA Group does not warrant the accuracy, completeness, or currency of any of the information published in this document. CSA Group makes no representations or warranties regarding this document’s compliance with any applicable statute, rule, or regulation.

IN NO EVENT SHALL CSA GROUP, ITS VOLUNTEERS, MEMBERS, SUBSIDIARIES, OR AFFILIATED COMPANIES, OR THEIR EMPLOYEES, DIRECTORS, OR OFFICERS, BE LIABLE FOR ANY DIRECT, INDIRECT, OR INCIDENTAL DAMAGES, INJURY, LOSS, COSTS, OR EXPENSES, HOWSOEVER CAUSED, INCLUDING BUT NOT LIMITED TO SPECIAL OR CONSEQUENTIAL DAMAGES, LOST REVENUE, BUSINESS INTERRUPTION, LOST OR DAMAGED DATA, OR ANY OTHER COMMERCIAL OR ECONOMIC LOSS, WHETHER BASED IN CONTRACT, TORT (INCLUDING NEGLIGENCE), OR ANY OTHER THEORY OF LIABILITY, ARISING OUT OF OR RESULTING FROM ACCESS TO OR POSSESSION OR USE OF THIS DOCUMENT, EVEN IF CSA GROUP HAS BEEN ADVISED OF THE POSSIBILITY OF SUCH DAMAGES, INJURY, LOSS, COSTS, OR EXPENSES.

In publishing and making this document available, CSA Group is not undertaking to render professional or other services for or on behalf of any person or entity or to perform any duty owed by any person or entity to another person or entity. The information in this document is directed to those who have the appropriate degree of experience to use and apply its contents, and CSA Group accepts no responsibility whatsoever arising in any way from any and all use of or reliance on the information contained in this document.

CSA Group is a private not-for-profit company that publishes voluntary standards and related documents. CSA Group has no power, nor does it undertake, to enforce compliance with the contents of the standards or other documents it publishes.

Intellectual property rights and ownership

As between CSA Group and the users of this document (whether it be in printed or electronic form), CSA Group is the owner, or the authorized licensee, of all works contained herein that are protected by copyright, all trade-marks (except as otherwise noted to the contrary), and all inventions and trade secrets that may be contained in this document, whether or not such inventions and trade secrets are protected by patents and applications for patents. Without limitation, the unauthorized use, modification, copying, or disclosure of this document may violate laws that protect CSA Group’s and/or others’ intellectual property and may give rise to a right in CSA Group and/or others to seek legal redress for such use, modification, copying, or disclosure. To the extent permitted by licence or by law, CSA Group reserves all intellectual property rights in this document.

Patent rights

Attention is drawn to the possibility that some of the elements of this standard may be the subject of patent rights. CSA Group shall not be held responsible for identifying any or all such patent rights. Users of this standard are expressly advised that determination of the validity of any such patent rights is entirely their own responsibility.

Authorized use of this document

This document is being provided by CSA Group for informational and non-commercial use only. The user of this document is authorized to do only the following:

If this document is in electronic form:

- load this document onto a computer for the sole purpose of reviewing it;
- search and browse this document; and
- print this document if it is in PDF format.

Limited copies of this document in print or paper form may be distributed only to persons who are authorized by CSA Group to have such copies, and only if this Legal Notice appears on each such copy.

In addition, users may not and may not permit others to

- alter this document in any way or remove this Legal Notice from the attached standard;
- sell this document without authorization from CSA Group; or
- make an electronic copy of this document.

If you do not agree with any of the terms and conditions contained in this Legal Notice, you may not load or use this document or make any copies of the contents hereof, and if you do make such copies, you are required to destroy them immediately. Use of this document constitutes your acceptance of the terms and conditions of this Legal Notice.



Revision History

CSA N288.1:20, Guidelines for modelling radionuclide environmental transport, fate, and exposure associated with the normal operation of nuclear facilities

Errata — November 2020	Revision symbol (in margin)
Tables C.5 , C.6 , G.2 , G.3 , G.4 , G.8 , and G.9c	Δ

Standards Update Service

CSA N288.1:20

March 2020

Title: *Guidelines for modelling radionuclide environmental transport, fate, and exposure associated with the normal operation of nuclear facilities*

To register for e-mail notification about any updates to this publication

- go to store.csagroup.org
- click on **Product Updates**

The **List ID** that you will need to register for updates to this publication is **2427651**.

If you require assistance, please e-mail techsupport@csagroup.org or call 416-747-2233.

Visit CSA Group's policy on privacy at www.csagroup.org/legal to find out how we protect your personal information.

CSA N288.1:20

***Guidelines for modelling
radionuclide environmental
transport, fate, and exposure
associated with the normal
operation of nuclear facilities***



®A trademark of the Canadian Standards Association, operating as "CSA Group"

*Published in March 2020 by CSA Group
A not-for-profit private sector organization
178 Rexdale Boulevard, Toronto, Ontario, Canada M9W 1R3*

*To purchase standards and related publications, visit our Online Store at store.csagroup.org
or call toll-free 1-800-463-6727 or 416-747-4044.*

ISBN 978-1-4883-2579-3

*© 2020 Canadian Standards Association
All rights reserved. No part of this publication may be reproduced in any form whatsoever
without the prior permission of the publisher.*

Contents

Technical Committee on Environmental Management for Nuclear Facilities	6
Subcommittee on Guidelines for Calculating Derived Release Limits for Radioactive Material in Airborne and Liquid Effluents for Normal Operation of Nuclear Facilities	9
Preface	11
0 Introduction	13
0.1 Application of the Guidelines	13
0.1.1 General	13
0.1.2 Derived release limit (DRL)	13
0.1.3 Dose to public	13
0.1.4 Environmental risk assessment (ERA)	14
0.1.5 Radionuclide relative contributions to dose	14
0.2 History of the Guidelines	14
1 Scope	15
1.1 Facilities	15
1.2 Release paths	15
1.3 Release duration	15
1.4 Contaminants	16
1.5 Receptors	16
1.6 Downwind distance of validity	16
1.7 Site specificity	16
1.8 Level of complexity	16
1.8.1 Simpler approaches	16
1.8.2 Default transfer parameters	16
1.9 Terminology	17
2 Reference publications	17
3 Definitions and abbreviations	29
3.1 Definitions	29
3.2 Abbreviations	32
4 Basic concepts	33
4.1 Dose limits	33
4.2 Representative person concept	34
4.2.1 Grouping members of the public	34
4.2.2 Age classes	34
4.2.3 Multiple representative persons	34
4.2.4 Indigenous and other representative persons	35
4.2.5 Homogeneity and group size	35
4.2.6 Averaging across distinct practices	35
4.2.7 Data for the representative person	36
4.2.8 Nuclear facility workers	36
4.2.9 Conservative dose estimates for the representative person	36

4.3	Selection of radionuclides	36
5	Environmental transfer model	45
5.1	General concepts	45
5.2	Calculation of derived release limits (DRLs)	51
5.2.1	General	51
5.2.2	DRLs for releases to air	52
5.2.3	DRLs for releases to surface water	52
5.2.4	Dose per unit release	52
5.2.5	Dose per unit release to air	52
5.2.6	Dose per unit release to surface water	52
5.2.7	Stochastic and deterministic effects and limits	53
5.2.8	General procedure for DRL calculation	53
6	Transfer parameters for airborne releases	54
6.1	Atmospheric dispersion (P_{01})	54
6.1.1	Air concentrations	54
6.1.2	Effective release height	55
6.1.3	Vertical dispersion and building wake effects	59
6.1.4	Meteorological data	61
6.1.5	Decay and ingrowth	63
6.1.6	Tritium transformation from HT to HTO	64
6.2	Air immersion ($P(e)_{19}$)	66
6.2.1	Semi-infinite cloud model	66
6.2.2	Finite cloud model	67
6.2.3	Dose coefficients	68
6.2.4	Occupancy factors	69
6.2.5	Building shielding factors	69
6.2.6	Noble gases	69
6.2.7	Radionuclides limited by dose to skin	70
6.2.8	Default values of $P(e)_{19}$	71
6.3	Radionuclide concentration in soil due to atmospheric deposition (P_{13})	71
6.3.1	Soil compartment	71
6.3.2	Soil model	72
6.3.3	Deposition velocity	73
6.3.4	Erosion losses	75
6.3.5	Volatilization losses	76
6.3.6	Losses due to leaching	76
6.3.7	Losses due to crop removal	77
6.4	Deposition on crops, forage, and wild plants (P_{14})	78
6.4.1	Deposition model	78
6.4.2	Values of f_{int}	78
6.4.3	Values of tf	78
6.4.4	Values of λ_p	79
6.4.5	Values of t_e	79
6.4.6	Tritiated water	79
6.4.7	Elemental tritium	79
6.4.8	Organically bound tritium	80
6.4.9	Carbon-14 (C-14)	81

6.5	Transfer from soil to groundwater wells ($P_{3area2w}$)	82
6.5.1	Conceptual well model	82
6.5.2	Groundwater travel time	82
6.5.3	Transfer to well water	84
6.5.4	Tritiated water	85
6.5.5	C-14	86
6.6	Transfer from soil to inland pond ($P_{3area2p}$)	86
6.6.1	Pond model	86
6.6.2	Parameter values for the pond model	87
6.7	Transfer from atmosphere to inland pond (P_{12p})	90
6.7.1	Pond model	90
6.7.2	Parameter values	90
6.7.3	Tritiated water	90
6.7.4	C-14	90
6.8	Plant uptake from soil (P_{3mass4})	91
6.9	Transfer from ponds and wells to animal and plant produce (P_{2p5} and P_{2w5} , P_{2p6} and P_{2p7})	92
6.9.1	Radionuclides other than tritium and noble gases	92
6.9.2	Tritiated water	93
6.9.3	Organically-bound tritium	94
6.9.4	P_{2p6} and P_{2p7} for tritium and C-14	95
6.9.5	Noble gases	95
6.10	Transfer from plants and invertebrates to animal produce (P_{45} , P_{65} , P_{75})	95
6.10.1	Radionuclides other than tritium, C-14, and noble gases	95
6.10.2	Tritiated water	96
6.10.3	Organically-bound tritium	97
6.10.4	C-14 and noble gases	98
6.10.5	Transfer to honey ($P_{45-honey}$)	99
6.11	Transfer from soil and sediment to animal produce (P_{3mass5} , P_{85})	100
6.11.1	Radionuclides other than tritium, C-14, and noble gases	100
6.11.2	Tritium, C-14, and noble gases	100
6.12	Transfer from air to animal produce (P_{15})	101
6.12.1	Radionuclides other than tritium, C-14, and noble gases	101
6.12.2	Tritium	101
6.12.3	C-14 and noble gases	102
6.13	Transfer from air to humans ($P(i)_{19}$)	102
6.14	External dose from ground deposits ($P(e)_{3area9}$)	103
6.15	Internal dose from ingestion ($P(i)_{29}$, $P(i)_{3mass9}$, P_{49} , and P_{59})	104
6.15.1	Human dietary intakes	104
6.15.2	Soil intake	105
6.15.3	Drinking water intake rates	105
6.15.4	Ingestion dose	106
6.16	Dose from water immersion ($P(e)_{2w9}$)	107
6.16.1	External dose	107
6.16.2	Internal dose from tritium	108
7	Transfer parameters for releases to surface water	109
7.1	Aquatic dispersion (P_{02})	109
7.1.1	Liquid emissions	109
7.1.2	Radionuclide concentrations in receiving waters	109

7.1.3	Discharges to coastal waters	110
7.1.4	Discharges to rivers	112
7.2	Contamination of soil by irrigation (P_{23})	114
7.2.1	Irrigation	114
7.2.2	Transfer from irrigation water to soil	115
7.2.3	Parameter values for the irrigation model	115
7.2.4	Re-suspension of soil (P_{3area1})	116
7.2.5	Volatilization from irrigated soil (P_{3mass1})	116
7.3	Direct contamination of crops by spray irrigation (P_{24})	118
7.3.1	Radionuclides other than tritium, C-14, and noble gases	118
7.3.2	Tritiated water (HTO)	118
7.3.3	Organically bound tritium (OBT)	119
7.3.4	C-14	119
7.4	Transfer from surface water to animal produce (P_{25})	120
7.5	Internal dose from drinking water ($P(i)_{29}$)	120
7.6	Dose from water immersion ($P(e)_{29}$)	120
7.6.1	External dose	120
7.6.2	Internal dose from tritium	121
7.7	Transfer to aquatic animals (P_{26}) and plants (P_{27})	122
7.7.1	Bioaccumulation factors	122
7.7.2	Fresh water	122
7.7.3	Sea water	123
7.7.4	Tritium	123
7.7.5	C-14	125
7.8	Transfer to sediment (P_{28})	126
7.9	Transfer from aquatic biota and sediment to animal produce	127
7.9.1	Transfer from aquatic animals (P_{65}) and plants (P_{75}) to animal produce	127
7.9.2	Transfer from sediment to animal produce (P_{85})	127
7.10	External dose from contaminated shoreline ($P(e)_{89}$)	128
7.11	Internal dose from incidental ingestion of contaminated sediments ($P(i)_{89}$)	128
7.12	Internal dose from intake of aquatic foods and irrigated crops	129

8 Application guides 130

8.1	Source averaging times for monitoring	130
8.1.1	Airborne releases	130
8.1.2	Liquid releases	131
8.2	Release duration	131
8.2.1	General considerations	131
8.2.2	Intermittent releases	131
8.2.3	Airborne releases	131
8.2.4	Liquid releases	132
8.2.5	Alternative approaches	133
8.3	Additivity effects from different radionuclides, pathways, and sources	133
8.4	Practical monitoring considerations	134

Annex A	— Default values of transfer parameters	136
Annex B	— Sample calculations	337
Annex C	— Dose coefficients	353
Annex D	— Selection of limiting radionuclides for mixed effluents	380

Annex E — Half-lives and decay constants for the radionuclides considered in these Guidelines	382
Annex F — Hydrologic and aquatic transport models	386
Annex G — Parameter values for terrestrial pathways models	393
Annex H — The finite cloud model for air immersion dose	412
Annex I — Equations for transfer factors for progeny in decay chains	466

Preface

This is the fourth edition of CSA N288.1, *Guidelines for modelling radionuclide environmental transport, fate and exposure associated with the normal operation of nuclear facilities*. It supersedes the previous editions published in 2014, 2008, and 1987 under the title, *Guidelines for calculating derived release limits for radioactive material in airborne and liquid effluents for normal operation of nuclear facilities*.

Major changes to this edition include

- a) updated the structure of these Guidelines for increased consistency within the CSA N288 series;
- b) removed administrative language from the Guidelines;
- c) provided additional clarification in areas identified by users;
- d) identified stable carbon data for aquatic plants and invertebrates;
- e) improved consistency of language throughout the Guidelines; and
- f) updated the guidance on transfer factors.

The COG background document (Hart, 2013) was revised in concert with the fourth edition; the updated version is referred to hereafter as the COG *Derived Release Limits Guidance* or CDG (Hart, 2019).

Users of these Guidelines are reminded that the site selection, design, manufacture, construction, installation, commissioning, operation, and decommissioning of nuclear facilities in Canada are subject to the *Nuclear Safety and Control Act* and its *Regulations*.

The CSA N-Series Standards provide an interlinked set of requirements for the management of nuclear facilities and activities. CSA N286 provides overall direction to management to develop and implement sound management practices and controls, while the other CSA Group nuclear Standards provide technical requirements and guidance that support the management system. These Guidelines work in harmony with CSA N286 and do not duplicate the generic requirements of CSA N286; however, they may provide more specific direction for those requirements.

These Guidelines were prepared by the Subcommittee on Guidelines for Calculating Derived Release Limits for Radioactive Material in Airborne and Liquid Effluents for Normal Operation of Nuclear Facilities, under the jurisdiction of the Technical Committee on Environmental Management and the Nuclear Strategic Steering Committee, and have been formally approved by the Technical Committee.

Notes:

- 1) *Use of the singular does not exclude the plural (and vice versa) when the sense allows.*
- 2) *Although the intended primary application of this Guideline is stated in its Scope, it is important to note that it remains the responsibility of the users of the Guideline to judge its suitability for their particular purpose.*
- 3) *This Guideline was developed by consensus, which is defined by CSA Policy governing standardization — Code of good practice for standardization as “substantial agreement. Consensus implies much more than a simple majority, but not necessarily unanimity”. It is consistent with this definition that a member may be included in the Technical Committee list and yet not be in full agreement with all clauses of this Guideline.*
- 4) *To submit a request for interpretation of this Guideline, please send the following information to inquiries@csagroup.org and include “Request for interpretation” in the subject line:*
 - a) *define the problem, making reference to the specific clause, and, where appropriate, include an illustrative sketch;*
 - b) *provide an explanation of circumstances surrounding the actual field condition; and*
 - c) *where possible, phrase the request in such a way that a specific “yes” or “no” answer will address the issue.*

Committee interpretations are processed in accordance with the CSA Directives and guidelines governing standardization and are available on the Current Standards Activities page at standardsactivities.csa.ca.

- 5) *This Guideline is subject to review within five years from the date of publication. Suggestions for its improvement will be referred to the appropriate committee. To submit a proposal for change, please send the following information to inquiries@csagroup.org and include "Proposal for change" in the subject line:*
- a) *Guideline designation (number);*
 - b) *relevant clause, table, and/or figure number;*
 - c) *wording of the proposed change; and*
 - d) *rationale for the change.*

CSA N288.1:20

Guidelines for modelling radionuclide environmental transport, fate, and exposure associated with the normal operation of nuclear facilities

0 Introduction

0.1 Application of the Guidelines

0.1.1 General

CSA N288.1 provides guidance for developing a radionuclide transport and exposure pathways model (referred to as "the model") which can be used in multiple applications including

- a) to calculate derived release limits for radionuclides corresponding to the regulatory annual dose limit for a member of the public;
- b) to calculate public dose based on radionuclide in effluent and/or environmental monitoring data; and
- c) to support environmental risk assessment for human health and ecological risk.

Note: *Environmental risk assessments (ERAs) are addressed in CSA N288.6.*

The Guidelines consist of an array of mathematical equations that describe the transfer of radioactive materials through the environment from either a point of release to a receptor or from an intermediate compartment to the receptor. It incorporates numerous transport and exposure pathways, which can lead to internal and/or external exposure resulting in a dose.

0.1.2 Derived release limit (DRL)

In the past, the Guidelines have most commonly been applied for the development of exposure-based radionuclide release rates, known as derived release limits or DRLs. The DRL for a given radionuclide is the release rate that would cause an individual of the most highly exposed group to receive a committed dose equal to the the annual public dose limit due to release of the radionuclide to air or surface water during normal operation of a nuclear facility over the period of a calendar year. The DRL is derived using mathematical equations that describe the transfer of radioactive materials through the environment to humans. It takes into account all exposure pathways, including external exposure from immersion in contaminated air and water, external exposure to contaminated substrate (sediment, soil, beach sand), and internal exposure from inhalation and ingestion of radioactivity.

Note: *For purposes of standardization, a DRL is based on 1 mSv/year, the dose limit for members of the public (see Clause 4.1). A facility's release limit approved by the authority having jurisdiction (AHJ), might be the DRL or a fraction of the DRL.*

0.1.3 Dose to public

These Guidelines can be applied to calculate the dose to a member of the public based on radionuclide release rates from a facility, which can be corroborated with radiological measurements in the environment where available.

0.1.4 Environmental risk assessment (ERA)

While these Guidelines focus on the calculation of dose to the public, they can also support the calculation of absorbed dose to non-human biota through the modelling of activity levels in environmental media and associated food webs. As for public dose, this modelling can be further corroborated with environmental measurements.

Note: An understanding of the fate and transport of contaminants in the environment is necessary for performing ERAs. The requirements for ERAs are addressed in CSA N288.6.

0.1.5 Radionuclide relative contributions to dose

When applied for either DRLs or for dose calculations, the model can provide information on the relative contribution to dose for each radionuclide and/or exposure pathway.

0.2 History of the Guidelines

In 1987, CSA Group published CSA N288.1 to provide guidelines and a methodology for calculating DRLs for routine releases of radionuclides to air and surface water from nuclear facilities. Since 1987, significant scientific advances were made in dosimetry and in the understanding of radionuclide behaviour in the environment; thus, there was a general consensus in the Canadian nuclear industry that the models and data in CSA N288.1 needed to be updated. Revision began in 2000 with the development of a derived release limits guidance document prepared for Ontario Power Generation (OPG). This document was specifically designed to calculate DRLs at CANDU® nuclear power generation facilities in southern Ontario (i.e., the Bruce, Pickering, and Darlington stations).

In 2006, the document was extended by the CANDU Owners Group (COG) to cover all CANDU facilities in Canada, including the power generation stations at Gentilly (G-2) and Point Lepreau and the research laboratories at Chalk River (CRL). This extended document (Hart, 2008) formed the basis for the second edition of these Guidelines and included background material deemed too detailed for inclusion in the Guidelines itself. Most of the models and parameter values in the first edition of the Guidelines were updated and new exposure pathways were added in producing the second edition, which was issued in 2008. Following publication, users of the Guidelines identified a number of minor errors, which were corrected in an amendment issued in July 2011.

It was not possible during the development of the second edition to implement all of the changes that were deemed desirable. These changes were set aside so that the second edition could be published in a timely fashion. Work on these changes continued through a number of research projects funded by COG. The findings of these projects form the basis for most of the improvements to the third edition, published in 2014. The COG background document (Hart, 2008) was revised in concert with the third edition (Hart, 2013).

Major updates to the third edition included the following:

- a) updated energy expenditures and dietary intake rates for humans;
- b) updated half-lives, gamma energies, and photon yields for all radionuclides;
- c) updated values for many parameters based largely on a new International Atomic Energy Agency handbook of parameter values for environmental transfers of radionuclides (IAEA, 2010);
- d) improved direction on when the Guidelines can be used to calculate DRLs for intermittent releases;
- e) updated wind direction and precipitation data for use in the wet deposition model;
- f) introduction of a model for wild waterfowl as an additional source of human exposure through ingestion;
- g) extension of the carbon-14 (C-14) specific activity model to cover plant to animal transfer;
- h) an improved specific activity model for tritium in animals, including an update and extension of the water intake source fractions for fresh and dry feed; and

- i) provision of equations for explicit accounting of decay and progeny ingrowth in all physical media, as an alternative to the use of progeny-inclusive dose coefficients.

Note: CANDU (CANada Deuterium Uranium) is a registered trademark of Atomic Energy of Canada Limited (AECL).

1 Scope

1.1 Facilities

These Guidelines and the CDG are intended to apply to CANDU nuclear power stations in Canada. However, the radionuclides and environmental pathways addressed make these Guidelines applicable to releases from many other nuclear facilities, including research reactors, radioisotope processing facilities, waste processing facilities such as incinerators, and power reactors other than those of CANDU design, subject to the limitations detailed in Clauses [1.2](#) to [1.8](#). Application to other types of facilities such as fuel fabrication plants and refineries is limited by the radionuclides considered here (see Clause [4.3](#)). These Guidelines may be adapted to cover part of the needs of such facilities, but additional models or methodologies might be necessary for other parts. However, neither the radionuclides nor the models included in these Guidelines are complete enough to cover releases from sources such as uranium mines and mills, or permanent geologic disposal facilities. In addition, the pathways are incomplete for any facilities where extensive modelling of groundwater pathways is required.

1.2 Release paths

These Guidelines cover releases to the atmosphere and to surface water (both fresh and marine). They do not address releases to groundwater, although transfers from other media to groundwater wells and ponds are considered. Direct gamma irradiation from radioactivity inside the facility is not modelled because it does not involve a release.

1.3 Release duration

The methods specified in these Guidelines are designed for routine, continuous, low-level emissions. They also apply to periodic, short-term releases (see Clause [8.2](#)), provided that

- a) the releases are controlled and associated with normal operations;
- b) the release rate is roughly the same from event to event;
- c) for atmospheric releases, the total release duration exceeds approximately 1000 h in the year; for aquatic releases, at least one or two releases occur in each month of the year; and
- d) the releases occur randomly over time.

Where the requirement of Item d) is not met but the releases are known to occur at a particular time of day or year, these Guidelines apply only if the air (water) concentrations are calculated using the meteorological (hydrological) data in effect for that time.

Notes:

- 1) *Where non-random releases are calculated using time-appropriate data, it might be possible to relax the conditions on the release frequencies.*
- 2) *Releases that do not meet these conditions can use another model, such as that specified in CSA N288.2 for atmospheric releases.*
- 3) *For some facilities, intermittent releases occur predictably as spikes on a continuous base release. Such releases can be considered part of routine emissions and included in the DRL without special treatment if the total activity released in intermittent form is less than approximately 30% of the total release from the facility. The 30% cut-off is considered a small fraction of the overall uncertainty of the DRL estimates.*